



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

New Alaskan Birds *

RICHARD C. MCGREGOR.

Leucosticte kadiaka sp. nov.

Sp. Char.—Similar to *Leucosticte griseonucha* but with smaller bill and smaller, weaker feet and claws.

Type—♂ ad., No. 3048. McGregor Coll., Karluk, Kadiak Island, Alaska; March 14, 1897; collected by Cloudsley Rutter.

Habitat—Kadiak Island, Alaska.

Specimens at hand are in such worn plumage that wing and tail measurements are worthless. In spring birds from Karluk the bill is dark horn color, while in November examples it is yellow with a dark tip. Measurements in hundredths of an inch are here given for three males from Amaknak and four males from Kadiak.

	Exp. Culmen	Bill from nostril	Middle toe with claw
<i>L. griseonucha.</i>	.54	.42	.97
<i>kadiaka.</i>	.48	.38	.88

Melospiza sanaka sp. nov.

Sp. Char.—Similar to *Melospiza cinerea* but bill longer and more slender, middle toe with claw shorter.

Type—No. 3361. McGregor Coll., Sanak Island, Alaska; June 13, 1894; collected by C. Littlejohn. Exposed culmen, .61; bill from nostril, .48; middle toe with claw, 1.02.

Habitat.—Sanak and Popoff Islands, Alaska.

The Sanak birds at hand are all in worn spring plumage so that they cannot be compared with my fall birds from Dutch Harbor which are in fresh plumage. Mr. Chase Littlejohn kindly let me use his series of skins from Sanak.

The following table of measurements shows the amount and character of difference in size of the two forms.

MEASUREMENTS OF *Melospiza cinerea* AND *M. sanaka*

	Sanak		Amaknak	
Five examples of each.	Average	Extremes	Average	Extremes
Exp. culmen ♂	.60	.58-.63	.57	.54-.58
Exp. culmen ♀	.59	.57-.63	.56	.54-.60
Bill from nos. ♂	.48	.47-.49	.44	.44-.45
Bill from nos. ♀	.48	.46-.51	.45	.43-.47
Mid. toe with claw ♂	1.03	1.01-1.06	1.07	1.06-1.08

* An author's edition of 100 copies was distributed Nov. 25, 1900.